


Trigonometric Identities:

$\tan \theta = \frac{\sin \theta}{\cos \theta}$	$\sin^2 \theta + \cos^2 \theta = 1$	$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$
$\cot \theta = \frac{\cos \theta}{\sin \theta}$	$\tan^2 \theta + 1 = \sec^2 \theta$	$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$
$\csc \theta = \frac{1}{\sin \theta}$	$\cot^2 \theta + 1 = \csc^2 \theta$	$\sin(-\theta) = -\sin \theta$
$\sec \theta = \frac{1}{\cos \theta}$	$\sin(2\theta) = 2 \sin \theta \cos \theta$	$\cos(-\theta) = \cos \theta$
$\cot \theta = \frac{1}{\tan \theta}$	$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$ $= 2 \cos^2 \theta - 1$ $= 1 - 2 \sin^2 \theta$	$\tan(-\theta) = -\tan \theta$

The Unit Circle:



Top 20 Integrals (1-10):

- 1) $\int a \, dx = ax + c$
- 2) $\int x^n \, dx = \frac{x^{n+1}}{n+1} + c \quad (n \neq -1)$
- 3) $\int \frac{1}{x} \, dx = \ln|x| + c$
- 4) $\int e^x \, dx = e^x + c$
- 5) $\int a^x \, dx = \frac{a^x}{\ln a} + c$
- 6) $\int \ln x \, dx = x \ln x - x + c$
- 7) $\int \sin x \, dx = -\cos x + c$
- 8) $\int \cos x \, dx = \sin x + c$
- 9) $\int \tan x \, dx = \ln|\sec x| + c$
- 10) $\int \cot x \, dx = \ln|\sin x| + c$

Top 20 Integrals (11-20):

- 11) $\int \sec x \, dx = \ln|\sec x + \tan x| + c$
- 12) $\int \csc x \, dx = -\ln|\csc x + \cot x| + c$
- 13) $\int \sec^2 x \, dx = \tan x + c$
- 14) $\int \sec x \tan x \, dx = \sec x + c$
- 15) $\int \csc^2 x \, dx = -\cot x + c$
- 16) $\int \csc x \cot x \, dx = -\csc x + c$
- 17) $\int \tan^2 x \, dx = \tan x - x + c$
- 18) $\int \frac{1}{a^2 + x^2} \, dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$
- 19) $\int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \sin^{-1} \frac{x}{a} + c$
- 20) $\int \frac{1}{x\sqrt{x^2 - a^2}} \, dx = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + c$

Properties of Definite Integrals:

$$\begin{aligned} \int_a^b f(x) \, dx &= - \int_b^a f(x) \, dx \\ \int_a^a f(x) \, dx &= 0 \\ \int_a^b k \cdot f(x) \, dx &= k \int_a^b f(x) \, dx \\ \int_a^b -f(x) \, dx &= - \int_a^b f(x) \, dx \\ \int_a^b (f(x) \pm g(x)) \, dx &= \int_a^b f(x) \, dx \pm \int_a^b g(x) \, dx \\ \int_a^c f(x) \, dx &= \int_a^b f(x) \, dx + \int_b^c f(x) \, dx \end{aligned}$$

Improper Integrals, $f(x)$ Continuous on $[a, \infty)$:

$$\int_a^{\infty} f(x) \, dx = \lim_{b \rightarrow \infty} \int_a^b f(x) \, dx$$

L'Hopital's Rule for Indeterminate Limits:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Exponential Growth/Decay:

$$\text{Differential Equation: } \frac{dy}{dt} = ky$$

$$\text{Exponential Growth Equation: } y = y_0 e^{kt}$$

Limit Definition of a Derivative

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Alternate Definition (at a Point)

$$f'(a) = \lim_{b \rightarrow a} \frac{f(b) - f(a)}{b - a}$$

Product Rule:

$$\frac{d}{dx}(fg) = f \frac{dg}{dx} + g \frac{df}{dx}$$

Quotient Rule:

$$\frac{d}{dx}\left(\frac{H_i}{L_o}\right) = \frac{L_o dH_i - H_i dL_o}{L_o L_o}$$

Chain Rule:

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x) \quad \text{OR} \quad \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

Power Rule:

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

Derivative of a Constant:

$$\frac{d}{dx}(c) = 0$$

Exponential/Logarithmic Derivatives:

$$\frac{d}{dx}(a^u) = \ln(a)a^u u'$$

$$\frac{d}{dx}(e^u) = \frac{du}{dx} e^u$$

Inverse Function:

$$\frac{d}{dx}(f^{-1}(x)) = \frac{1}{f'(f^{-1}(x))}$$

Intermediate Value Theorem:

If a and b are any two points in an interval on which f is continuous, then f takes on every value between $f(a)$ and $f(b)$.

Intermediate Value Theorem for Derivatives: If a and b are any two points in an interval on which f is differentiable, then the derivative f' takes on every value between $f'(a)$ and $f'(b)$.

Average Rate of Change of a function f on $[a,b]$

$$\frac{f(b) - f(a)}{b - a}$$

Instantaneous Rate of Change of a function of f at $x = a$:

$$f'(c) = \frac{f(b) - f(a)}{b - a} \quad (\text{at some point "IROC" = "AROC"})$$

Average Value of a Function on an Interval $[a,b]$:

$$\text{Average Value of } f(x) \text{ on } [a,b] = \frac{1}{b-a} \int_a^b f(x) \, dx$$

Mean Value Theorem (for Definite Integrals):

If f is continuous on $[a,b]$, then at some point c in (a,b) such that

$$f(c) = \frac{1}{b-a} \int_a^b f(x) \, dx$$

(At some point, the function equals its average value on the interval.)

Fundamental Theorem of Calculus (Part 1):

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x) \quad \text{OR} \quad \frac{d}{dx} \int_a^u f(t) \, dt = u' f(u)$$

Fundamental Theorem of Calculus (Part 2):

$$\int_a^b f(x) \, dx = F(b) - F(a) \quad [F(x) \text{ is an antiderivative of } f(x)]$$

Integration by Parts:

$$\int u \, dv = uv - \int v \, du$$

$$\int f(x)g'(x) \, dx = f(x)g(x) - \int f'(x)g(x) \, dx$$

Logistic Growth/Decay:

$$\text{Differential Equation: } \frac{dP}{dt} = kP(M-P)$$

$$\text{Logistic Equation: } P(t) = \frac{M}{1 + Ae^{-(Mt)}} \quad (k \text{ is the growth/decay constant})$$

(M is the carrying capacity)

(A is a constant you must solve for)

-The population is growing the fastest when P is half of the carrying capacity!

-As t tends to infinity, the population tends to the carrying capacity:

$$\lim_{t \rightarrow \infty} P(t) = M$$

Volume (Discs):

$$V_{\text{discs about } x\text{-axis}} = \pi \int_a^b f(x)^2 dx$$

$$V_{\text{discs about } y\text{-axis}} = \pi \int_c^d f(y)^2 dy$$

Volume (Shells):

$$V_{\text{shells about } x\text{-axis}} = 2\pi \int_c^d y f(y) dy$$

$$V_{\text{shells about } y\text{-axis}} = 2\pi \int_a^b x f(x) dx$$

Volume (Cross Sections):

$$V_{\text{cross sections } \perp x\text{-axis}} = \int_a^b A(x) dx$$

$$V_{\text{cross sections } \perp y\text{-axis}} = \int_c^d A(y) dy$$

Arc Length (y a function of x):

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Arc Length (x a function of y):

$$L = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

Arc Length (Parameterized Curve):

$$L = \int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Surface Area (Rotate About x-axis):

$$SA = 2\pi \int_a^b f(x) \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Surface Area (Rotate About y-axis):

$$SA = 2\pi \int_c^d f(y) \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

Surface Area (Rotate Parameterized Curve About The x-axis):

$$SA = 2\pi \int_{t_1}^{t_2} y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Position, Velocity, and Acceleration Vectors:

$$\mathbf{r}(t) = \langle x(t), y(t) \rangle$$

$$\mathbf{v}(t) = \langle x'(t), y'(t) \rangle$$

$$\mathbf{a}(t) = \langle x''(t), y''(t) \rangle$$

Speed at time t_0 :

$$|v(t_0)| = \sqrt{x'(t_0)^2 + y'(t_0)^2}$$

Total Distance, $t \in [a,b]$:

$$\begin{aligned} \text{Total Distance} &= \int_a^b |v(t)| dt \\ &= \int_a^b \sqrt{(v_1(t))^2 + (v_2(t))^2} dt \end{aligned}$$

Displacement, $t \in [a,b]$:

$$\left\langle \int_a^b v_1(t) dt, \int_a^b v_2(t) dt \right\rangle$$

Final Position, $t \in [a,b]$:

$$\langle x(a), y(a) \rangle + \left\langle \int_a^b v_1(t) dt, \int_a^b v_2(t) dt \right\rangle$$

Linearization:

If f is differentiable at $x = a$, then the equation of the tangent line,

$L(x) = f(a) + f'(a)(x - a)$, defines the linearization of f at a . The standard linear approximation of f at a is $f(x) \approx L(x)$. The point $x = a$ is the center of the approximation. This is just a Degree 1 Taylor Series approximation of f at a !!

Newton's Method for Approximating a Solution to $f(x) = 0$:

1. Guess a first approximation to a solution of the equation $f(x) = 0$.
2. Use the first approximation to get a second, the second to get a third, and so on, using the formula:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Euler's Method for Approximating $f(a)$:

1. Start at a point (x, y) specified by an initial condition.
2. Use the differential equation to find the slope dy/dx at (x, y)
3. Move by a small increment, Δx , and use this to determine Δy using $\Delta y = (dy/dx)\Delta x$.
4. Use the new point, $(x + \Delta x, y + \Delta y)$, then repeat from Step 2.
5. Continue until you have your approximation.

(x, y)	dy/dx	Δx	$\Delta y = (dy/dx)\Delta x$	$(x + \Delta x, y + \Delta y)$

Geometric Series:

$$\sum_{n=1}^{\infty} a_n r^{n-1} = a_1 + a_2 r + a_3 r^2 + \dots + a_n r^{n-1} + \dots = \frac{a_1}{1-r} \text{ for } |r| < 1$$

Taylor Series for $f(x)$ centered at $x = a$:

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n$$

MacLaurin for $f(x)$ (always centered at $x = 0$):

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}x^n$$

Actual Error (Actual Minus Estimate):

$$|R_n(x)| = |f(x) - P_n(x)|$$

LaGrange Error Bound on $[a, b]$:

$$|R_n(x)| < \frac{|f^{(n+1)}(c)x^{n+1}|}{(n+1)!} \quad (\text{Pick } c \text{ to maximize } f^{(n+1)}(c))$$

Alternating Series:

The error is no more than the next term!

MacLaurin Series To Memorize (Part 1):

$$\frac{1}{1-x} = 1 + x + x^2 + \dots + x^n + \dots \quad (|x| < 1)$$

$$\frac{1}{1+x} = 1 - x + x^2 - \dots + (-x)^n + \dots \quad (|x| < 1)$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (\text{for all real } x)$$

MacLaurin Series To Memorize (Part 2):

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \quad (\text{for all real } x)$$

$$\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^n \frac{x^{2n}}{(2n)!} + \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad (\text{for all real } x)$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^{n-1} \frac{x^n}{n} + \dots = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} \quad (-1 < x \leq 1)$$

$$\tan^{-1}(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)} + \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)} \quad (|x| \leq 1)$$

Convergence at Endpoints: When you have an infinite series involving x , use Ratio Test to find an open interval of convergence. Then use other tests at endpoints!

Ratio Test:

$$\text{For } \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L :$$

$L < 1$: the series converges.

$L > 1$: the series diverges.

$L = 1$: test is inconclusive.

Limit Comparison:

If $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$ and $\sum b_n$ converges, then so does $\sum a_n$.

If $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$, then both converge or diverge.

Alternating Series

- 1) Show terms alternate
- 2) Show $a_n \geq a_{n+1}$
- 3) Show $\lim_{n \rightarrow \infty} |a_n| = 0$

If so, then the series converges.

Polar Coordinates:

$$\begin{aligned} x &= r \cos \theta & y &= r \sin \theta & r^2 &= x^2 + y^2 & \tan \theta &= \frac{y}{x} \end{aligned}$$

Parameterize the Polar Equation $r = f(\theta)$:

$$x(\theta) = f(\theta) \cos(\theta)$$

$$y(\theta) = f(\theta) \sin(\theta)$$

Finding dy/dx (Slope) for a Polar Curve:

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} \quad \left(\frac{dx}{d\theta} \neq 0 \right)$$

Finding Area for a Polar Curve:

$$A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta = \frac{1}{2} \int_{\alpha}^{\beta} f(\theta)^2 d\theta$$

p-series Test:

$$\sum \frac{1}{n^p} \text{ converges if } p > 1$$

Geometric Series:

$$\sum_{n=1}^{\infty} a_n r^{n-1} \text{ converges if } |r| < 1$$

nth Term Test:

If $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series diverges.

Integral Test:
If $f(n) = a_n$ is a decreasing sequence, then $\sum a_n$ and $\int f(x) dx$ both converge or diverge.

Telescoping Series:
Use partial fraction decomposition to separate into two sequences, then group terms and cancel!